

Agromet Advisory Bulletin for the District, Kozhikode (Valid from 16.0.2025 to 20.07.2025)



(Issued jointly by Kerala Agricultural University Regional Agricultural Research Station Pilicode& India Meteorological Department)

Bulletin Number:Pilicode/ Kkd -56/2025 Date: 15/07/2025

A. Weather Summary of preceding five days

Rainfall, mm	Max. temp., ⁰C	Min. temp., °C	R. H., %	Wind speed, Km/h
90.1	28.9 - 32.5	23.1 - 25.0	76 – 96	00 - 00

B.Weather forecast for next five days

Parameters	16-07-2025	17-07-2025	18-07-2025	19-07-2025	20-07-2025
Average Rainfall, mm	20	25	30	20	18
Max. Temp, °C	29	29	29	29	29
Min. Temp,°C	23	23	23	23	23
Max. Relative Humidity, %	95	95	95	95	95
Min. Relative Humidity, %	90	90	90	90	90
Wind speed,km/h	3	3	3	3	4
Wind direction, degrees	320	340	320	320	340
Total cloud cover, octa	8	8	8	8	8

C. Agrometeorological Advisories

Сгор	Stages	Problems	Agro-meteorological advisories				
	Isolated Heavy To Very Rainfall**						
General Condition	The sky will be cloudy. High relative humidity will be experienced. The atmospheric temperature will come down. There will be heavy rainfalls on July16 in isolated places. There will be heavy to very heavy rainfalls from July17 to 19 in isolated places.						
Weather warning	Chances for isolated heavy rainfall on July15.						
Impacts	Difficulty in drying agricultural produce in open conditions.						
	Chances for fall off of slender stemmed fruits and vegetables.						
	Chances for water stagnation in the fields.						
General Recommendati ons	Drain the stagnating water from the cultivated areas where heavy water stagnations occurred due to the last five days heavy rainfalls						
0113	Harvest all the ripened products if any quickly. Do not keep the harvested products in the field itself.						
	Do not attempt to dry the products like rice, rubber, copra, pulses, vegetable seeds, etc direct under sun.						

	 Give popping to all soft slender stemmed crops like banana, vegetables, climbers etc. Clean the drainage channels in crop lowlands to enable proper drainage of excess water in case of exigencies if any. Do not shelter the cattle in houses with weak structures. Keep animal feeds, seeds, fertilizers etc. in termite free and moist free rooms on elevated platforms coated with wooden panels. 					
	 Avoid activities under open conditions while there is lightning. Also do not expose livestock to such unfavorable weather conditions. Seedlings of tree crops can be planted now. Ensure proper distance between the plants. The ²/₃ portions of the pits should be filled with mixture of topsoil and powdered dry cow dung/compost. Spraying should be done only if sufficient intervals are available between the rainfalls. Use sticking agents like 					
Paddy	Seedlings(Nursery/ Transplanted)	Leaf folder	If leaf folder attack is seen, unfold the leaves by rolling thorny wires/twigs over the canopy. If the attack is severe spot spraying of flubendiamide @ 1.0 ml/10L is recommended. Attack will be more in shaded conditions. Avoid excessive Nitrogenous fertilizers			
Coconut	Newly planted seedlings	Water stagnation and decaying of growing bud (bud rot)	Do not allow water stagnation in the planted pits. Provide bunds around the pits to prevent the collection of runoff water in the pits. After the cessation of heavy rainfall, spray 1 % Bordeaux Mixture on the leaves and growing buds, only after mixing with a sticking agent to the fungicide.			
Coconut	All stages	Rhinoceros beetle Warf varieties and young palms are more vulnerable to the attack.	Take out the beetles from the attacked palm crown using beetle hook. Swab the hole made by the beetles with Bordeaux paste to prevent entry of fungus through the cut surface. Old fishnets can be reused for controlling the pest. Cut the nets into pieces of size 1m x 0.5m. Keep the middle portion of the net, lengthwise into the axils of 3 rd and 4 th leaf whirls of the palm. Push the net firmly with small stones into the axils. The hanging sides of the net should be kept open. The beetles will be trapped in the nets and get destroyed. Care should be taken to move the nets to upper leaf axils accordingly while fresh leaves are emerging.			

Black pepper	Bearing stage	Pollu beetle	Mix quinalphos 25%EC (@ 2ml per litre) with 1% of Bordeaux mixture and spray
Black pepper	All stages	Foot rot	As prophylactic measure, apply 150 gram of Trichoderma enriched neem cake - cow dung mixture in the basins of the vines and incorporate thoroughly with the soil. If disease already appeared, drench soil in the plant basins with Redomil 0.2% (2g/litre of water). Spray the same on the leaves also.
Banana	Various stages of growth	Rhizome rot	Ensure good drainage in the garden Drench with Streprocyclin antibiotic (@2g/10L) Or Copperhydroxide (20g/10L) at the basin of the plant.
Poultry	Various stages of growth	Cage hygiene	Keep the cages dry. If moisture persists, ammonia may be generated from the droppings which will adversely affect the health of the birds.

**	Warning	colour	codes	of	rainfall	(for	disaster	management	t)
----	---------	--------	-------	----	----------	------	----------	------------	----

ſ	Warning (Take actions)	Alert (Be prepared)	Watch (Be updated)	No warning (No actions)		
L						

Sd/-Nodal Officer, GKMS Project, RARS Pilicode